

AD-A264 931 ITATION PAGE

Form Approved  
OBM No. 0704-0188

age 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and  
ormation. Send comments regarding this burden or any other aspect of this collection of information, including suggestions  
for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to  
0188), Washington, DC 20503.

June 1992

3. Report Type and Dates Covered.  
Final - Proceedings

## 4. Title and Subtitle.

Atmospheric Correction of AVIRIS Data in Ocean Waters

## 5. Funding Numbers.

Contract

Program Element No. 0601013N

Project No. R1947

Task No.

Accession No. DN251095

Work Unit No. 93502G

## 6. Author(s).

Gregory Terrie and Robert Arnone

## 7. Performing Organization Name(s) and Address(es).

Naval Research Laboratory  
Ocean Science Directorate  
Stennis Space Center, MS 39529-50048. Performing Organization  
Report Number.

PR 92:075:352

## 9. Sponsoring/Monitoring Agency Name(s) and Address(es).

Navy Engineering Logistics Office  
P.O. Box 15467  
Arlington, VA10. Sponsoring/Monitoring Agency  
Report Number.

PR 92:075:352

## 11. Supplementary Notes.

Published Jet Propulsion Laboratory Publication 92-14.

## 12a. Distribution/Availability Statement.

Approved for public release; distribution is unlimited.

## 12b. Distribution Code.

## 13. Abstract (Maximum 200 words).

Hyperspectral data offers unique capabilities for characterizing the ocean environment. The spectral characterization of the composition of ocean waters can be organized into biological and terrigenous components. Biological photosynthetic pigments in ocean waters have unique spectral ocean color signatures which can be associated with different biological species (Arnone et al., 1986). Additionally, suspended sediment has different scattering coefficients which result in ocean color signatures. Measuring the spatial distributions of these components in the maritime environments provides important tools for understanding and monitoring the ocean environment. These tools have significant applications in pollution, carbon cycle, current and water mass detection, location of fronts and eddies, sewage discharge and fate etc.

8 5 24 0 8

93-11541  
 SPJ

## 14. Subject Terms.

Coastal Zone Color Scanner, airborne visible-infrared imaging spectrometer

## 15. Number of Pages.

2

## 16. Price Code.

17. Security Classification  
of Report.

Unclassified

18. Security Classification  
of This Page.

Unclassified

19. Security Classification  
of Abstract.

Unclassified

## 20. Limitation of Abstract.

SAR

JPL Publication 92-14, Vol. 1

# Summaries of the Third Annual JPL Airborne Geoscience Workshop June 1-5, 1992

Volume 1. AVIRIS Workshop

Robert O. Green  
Editor

June 1, 1992



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

DTIC QUALITY INSPECTED 3

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

# ATMOSPHERIC CORRECTION OF AVIRIS DATA IN OCEAN WATERS

Gregory Terric and Robert Arnone  
Naval Research Laboratory, Stennis Space Center

## 1. INTRODUCTION

Hyperspectral data offers unique capabilities for characterizing the ocean environment. The spectral characterization of the composition of ocean waters can be organized into biological and terrigenous components. Biological photosynthetic pigments in ocean waters have unique spectral ocean color signatures which can be associated with different biological species (Arnone et al, 1986). Additionally, suspended sediment has different scattering coefficients which result in ocean color signatures. Measuring the spatial distributions of these components in the maritime environments provides important tools for understanding and monitoring the ocean environment. These tools have significant applications in pollution, carbon cycle, current and water mass detection, location of fronts and eddies, sewage discharge and fate etc.

Ocean color has been used from satellite for describing the spatial variability of chlorophyll, water clarity ( $k_{490}$ ), suspended sediment concentration, currents etc. Additionally, with improved atmospheric correction methods, ocean color results have produced global products of spectral water leaving radiance ( $L_w$ ) (Feldman 1989). Ocean color results have clearly indicated strong applications for characterizing the spatial and temporal variability of bio-optical oceanography (Arnone et al, 1991).

These studies have been largely the results of advanced atmospheric correction techniques applied to multispectral imagery. The atmosphere contributes approximately 80% - 90% of the satellite received radiance in the blue-green portion of the spectrum. In deep ocean waters, maximum transmission of visible radiance is achieved at 490nm. Conversely, nearly all of the light is absorbed by the water at wavelengths greater than about 650nm and thus appears black. These spectral ocean properties are exploited by algorithms developed for the atmospheric correction used in satellite ocean color processing.

The objective of this study was to apply atmospheric correction techniques that were used for processing satellite Coastal Zone Color Scanner (CZCS) data to AVIRIS data. Quantitative measures of  $L_w$  from AVIRIS will be compared with ship ground truth data and input into bio-optical models.

## 2. METHODS

Atmospheric correction of CZCS assumes a linear summation of the total radiance,  $L_t(\lambda)$ , sensed at the sensor (Gordon et al, 1983, Gordon and Clark, 1979). The atmospheric contribution is divided into the Rayleigh,  $L_r(\lambda)$ , component (molecular scattering) and the aerosol,  $L_a(\lambda)$ , component (Mie scattering). The volume scattering of the water,  $L_w(\lambda)$  (ocean color) is transmitted through the air sea interface,  $t(\lambda)$ . All of these parameters are related to the total radiance,  $L_t(\lambda)$  by

$$L_t(\lambda) = L_r(\lambda) + L_a(\lambda) + t(\lambda)L_w(\lambda). \quad (1)$$

Assuming that the signal at 670nm is due entirely to atmospheric effects and the aerosol optical thickness  $\tau_a(670)$  is proportional to  $\tau_a(\lambda)$ , equation (1) can be written in terms of  $L_w(\lambda)$ , at nadir, by

$$iL_w(\lambda) = L_i(\lambda) - L_r(\lambda) - \frac{F_0(\lambda)}{F_0(670)} \left( \frac{\lambda}{670} \right)^n (L_i(670) - L_r(670)), \quad (2)$$

where  $F_0$  is the solar flux and  $n$  is the angstrom coefficient.

This relationship was applied to AVIRIS data by expanding this generalized nadir equation to account for the angular atmospheric path radiance from the sensor altitude. The rayleigh component  $L_r(\lambda)$  was computed using a multiple scattering model (Gordon et al, 1988). Both solar and sensor zenith and azimuth angles were computed for each pixel in the scene according to the date, time and location of data collection. The solar fluxes were assumed constants and the angstrom coefficient was varied.

Four spectral channels were used in processing CZCS ocean color data which were centered at 443nm, 520nm, 550nm, and 670nm with a bandwidth of 20 nm. The 443nm channel represents the chlorophyll absorption wavelength which is inversely proportional to concentration (Clark 1980). The 520nm and 550nm channels characterize the hinge point in bio-optical ocean waters and are related to sediment scattering. These bands were simulated using AVIRIS bands by averaging the two adjacent bands for each wavelength. Additionally, a 750nm channel, analogous to the CZCS 750nm band, was used as the atmospheric reference channel because of a problem with bottom reflectance at 670nm, especially in the shallow coastal regions. Each of the channels were calibrated for spectral radiance and for vignetting effects using the supplied correction data.

The atmospheric model was applied to an AVIRIS scene collected off Tampa Bay on April 3, 1990. This image has many different ocean regimes characterizing, shallow coastal waters with a uniform bottom reflectance, suspended sediment in the water, absorbing "yellow substance", and variability of coastal phytoplankton and clear waters. Ship measurements were collected by the University of South Florida (Carder et al, 1991) coincident with the overflight at three stations in which  $L_w(\lambda)$ , chlorophyll, and the diffuse attenuation coefficient were measured.

### 3. RESULTS

The 443nm channel is severely contaminated by the atmosphere and sensor noise such that little ocean features are observed. Atmospheric scattering diminishes in the 520nm and 550nm channels and ocean features can be recognized although not quantitative. Atmospheric correction using the 670nm channel as the reference was applied to the scene ranging from clear waters to the coastal environments. In the near shore coastal waters, the AVIRIS data was shown to be correlated with the bathymetry since the reflectivity of the white bottom clearly dominated the water leaving radiance signal. The 670nm channel was shown to be correlated with bottom reflectivity and therefore is not a good channel for only characterizing the atmosphere. Because of these non zero 670nm components, there is an overcorrection for the atmosphere. This results in negative radiance in areas where the bottom reflectance dominates the 670nm channel. Since this is not possible the images were processed and negative values clipped to zero. The

Assuming that the signal at 670nm is due entirely to atmospheric effects and the aerosol optical thickness  $\tau_a(670)$  is proportional to  $\tau_a(\lambda)$ , equation (1) can be written in terms of  $L_w(\lambda)$ , at nadir, by

$$tL_w(\lambda) = L_r(\lambda) - L_r(\lambda) - \frac{F_0(\lambda)}{F_0(670)} \left( \frac{\lambda}{670} \right)^n (L_r(670) - L_r(670)), \quad (2)$$

where  $F_0$  is the solar flux and  $n$  is the angstrom coefficient.

This relationship was applied to AVIRIS data by expanding this generalized nadir equation to account for the angular atmospheric path radiance from the sensor altitude. The rayleigh component  $L_r(\lambda)$  was computed using a multiple scattering model (Gordon et al, 1988). Both solar and sensor zenith and azimuth angles were computed for each pixel in the scene according to the date, time and location of data collection. The solar fluxes were assumed constants and the angstrom coefficient was varied.

Four spectral channels were used in processing CZCS ocean color data which were centered at 443nm, 520nm, 550nm, and 670nm with a bandwidth of 20 nm. The 443nm channel represents the chlorophyll absorption wavelength which is inversely proportional to concentration (Clark 1980). The 520nm and 550nm channels characterize the hinge point in bio-optical ocean waters and are related to sediment scattering. These bands were simulated using AVIRIS bands by averaging the two adjacent bands for each wavelength. Additionally, a 750nm channel, analogous to the CZCS 750nm band, was used as the atmospheric reference channel because of a problem with bottom reflectance at 670nm, especially in the shallow coastal regions. Each of the channels were calibrated for spectral radiance and for vignetting effects using the supplied correction data.

The atmospheric model was applied to an AVIRIS scene collected off Tampa Bay on April 3, 1990. This image has many different ocean regimes characterizing, shallow coastal waters with a uniform bottom reflectance, suspended sediment in the water, absorbing "yellow substance", and variability of coastal phytoplankton and clear waters. Ship measurements were collected by the University of South Florida (Carder et al, 1991) coincident with the overflight at three stations in which  $L_w(\lambda)$ , chlorophyll, and the diffuse attenuation coefficient were measured.

### 3. RESULTS

The 443nm channel is severely contaminated by the atmosphere and sensor noise such that little ocean features are observed. Atmospheric scattering diminishes in the 520nm and 550nm channels and ocean features can be recognized although not quantitative. Atmospheric correction using the 670nm channel as the reference was applied to the scene ranging from clear waters to the coastal environments. In the near shore coastal waters, the AVIRIS data was shown to be correlated with the bathymetry since the reflectivity of the white bottom clearly dominated the water leaving radiance signal. The 670nm channel was shown to be correlated with bottom reflectivity and therefore is not a good channel for only characterizing the atmosphere. Because of these non-zero 670nm components, there is an overcorrection for the atmosphere. This results in negative radiance in areas where the bottom reflectance dominates the 670nm channel. Since this is not possible the images were processed and negative values clipped to zero. The